



# The Veterinary Link

by **BOB LARSON, DVM**, *University of Missouri-Columbia*

## Vaccination strategies to reduce pregnancy loss

The first step in designing a program to minimize pregnancy wastage (infertility, abortions and stillbirths) is to select the diseases against which the herd should be immunized.

Commercial vaccines are not available for all pregnancy-wasting diseases. Other diseases have vaccines manufactured for their control, but the vaccines may not be adequately effective or a primary concern for a particular area or herd. For most beef herds the potential list of diseases to include in a vaccination program would include brucellosis (Bang's disease), vibriosis (campylobacteriosis), leptospirosis (lepto), infectious bovine rhinotracheitis (IBR) and bovine viral diarrhea (BVD).

Other diseases for which vaccines are available include neosporosis, anaplasmosis, *Haemophilus somnus* and trichomoniasis (trich). When selecting diseases to protect against in a vaccination program, it is wasteful and unjustified to vaccinate with every available vaccine. The diseases selected should be limited to those for which

there are effective vaccines and to which the herd will possibly be exposed. This decision is largely based on the classification of the herd as a closed, modified-open or open herd.

A closed herd has no new additions except for bulls, and there is no animal-to-animal contact with neighboring herds. New bulls are isolated prior to introduction into the herd, where they are appropriately tested for and vaccinated against contagious diseases during the isolation period. The level of immunization necessary for a closed herd is less than that for other herds because the risk of exposure is very low.

Modified-open herds have a higher risk of exposure to pregnancy-wasting diseases through limited additions of bulls and replacement females, animals moving in and out of the herd to exhibitions, or through contact with neighboring herds.

Open herds have a very high risk of exposure to pregnancy-wasting pathogens. This exposure may be due to the frequent introduction of replacements or exposure of the

breeding herd to stocker cattle or other recently purchased, stressed cattle.

### Brucellosis

Since brucellosis is a disease that can be spread to humans, its control in animals is especially important. *Brucella abortus* strain RB51 vaccine is a live bacterial product and confers long-term cell-mediated protection in healthy animals vaccinated properly and produces fewer false-positive blood tests in vaccinated cattle than the older, strain 19 vaccine.

Official calftag vaccination of females between 4 and 12 months of age (some state requirements vary) is undertaken by accredited veterinarians or state or federal animal health representatives only. Whole-herd (adult) vaccination can only be undertaken under specific

*(Continued on page 42)*



[PHOTO BY LISA KLOENTRUP]

---

**Protection from pregnancy wastage in beef cattle has been enhanced by the development and improvement of vaccines against several important organisms causing embryonic loss and abortion.**

---

**The Veterinary Link** *(from page 41)*

situations with approval from a state or federal animal health official.

Brucellosis is also controlled by a monitoring system of blood testing and identifying market cattle, cattle at livestock sales and cattle moving across state lines. Positive test reactors and the herds in contact with positive reactors

are handled by rules developed by the U.S. Department of Agriculture (USDA). For most areas of the country, brucellosis has been eradicated, and many herds are no longer utilizing official calfhood vaccination. Whether or not to continue with a brucellosis vaccination program should be determined only after considering interstate movement, risk of exposure and legal responsibility.

**IBR and BVD**

IBR is a herpes virus infection and is easily transmitted by secretions from the respiratory tract, the eyes and the reproductive system. Like other herpes viruses, IBR viruses often cause hidden infections. The carriers that do not appear sick are important sources of exposure.

The BVD virus is widely distributed and is easily spread from one animal to

another. Infection with BVD causes different reproductive losses, depending on the time a susceptible female is exposed to the virus. Abortion losses to BVD are sporadic, and the rate is usually low. Frequently, BVD is found in fetuses aborted due to other agents, such as fungal organisms, because BVD may suppress a herd's immunity and increase susceptibility to other infections.

To decrease the risk of pregnancy wastage from these viral diseases, nonpregnant heifers should be given modified-live vaccines (MLV) two or more times from weaning to 6 weeks before breeding. Although modified-live IBR/BVD vaccines do not require a booster to induce a protective response, it is recommended that vaccinations be repeated two or more times. Repeated vaccination is needed because one does not know when maternal antibody interference with active immunization wanes or if nutritional or host factors interfering with immunization are present. Multiple vaccinations allow the maximum number of heifers to develop active immunity to the vaccination. An open herd with a high level of risk may benefit from having annual IBR/BVD booster immunizations.

Veterinarians and immunologists have not been able to agree on the superiority of modified-live BVD vaccines vs. killed BVD vaccines for immunization of heifers. But, all veterinarians agree that because of the complexity of BVD infection and disease, a well-thought-out plan that includes screening for persistently infected (PI) individuals, herd biosecurity and vaccination is necessary to protect against this persistently troublesome virus.

**Leptospirosis**

This bacterial disease causes pregnancy wastage primarily in the last trimester of gestation of cattle, as well as infertility in the first trimester. Leptospiral organisms cause infection in the kidneys and are excreted in urine. Leptospire survive in wet environments for up to 30 days. Infection of susceptible cattle occurs through mucous membranes and abraded or water-softened skin, or by sexual contact. There are more than 180 serovars of leptospira.

Each serovar is adapted to a particular species of animal, which it uses as a maintenance host. In the United States, serovar hardjo has a maintenance host relationship with cattle. A maintenance host relationship is characterized by high susceptibility to infection, easy transmission within the host species, relatively mild disease in a host, a tendency to cause long-term rather than sudden disease and poor effectiveness of vaccination for prevention. Infertility may follow localization of leptospire in the uterus and oviduct of maintenance host hardjo carriers. A newly available vaccination against hardjo infection in cattle appears to prevent kidney

*(Continued on page 47)*

## The Veterinary Link (from page 42)

establishment, urinary shedding and fetal infection.

By contrast, an incidental host relationship is characterized by relatively low susceptibility to infection, with a tendency to cause acute, severe disease. One also finds sporadic transmission within the incidental host species and good efficacy of vaccination for preventing infection. Serovar pomona is a common incidental pathogen of cattle, and the maintenance host is swine. *Leptospira* strains maintained by nondomestic animals such as skunks, raccoons, opossums, foxes, beavers, mice, deer and others can infect cattle herds that are exposed to environments such as urine-contaminated waterholes.

To establish immunity against the organism, primary immunization of heifers should consist of two or three vaccinations given at monthly intervals prebreeding, and another booster in midgestation of the first pregnancy. Annual or biannual boosters, at prebreeding and midgestation, should be given.

Methods other than vaccinations for reducing risk of exposure to leptospirosis should also be implemented. Options include having a closed herd and fencing cattle away from water sources that can be contaminated by other herds, swine or nondomestic animals.

### Vibriosis

*Campylobacter fetus* subspecies *venerealis* is a disease passed during the act of mating. Infection of the uterus and oviducts can persist for up to two months, but thereafter the organism is progressively eliminated.

Management factors that minimize risk include utilizing artificial insemination (AI) with semen from noninfected bulls; utilizing bulls less than 3 years of age, since they tend to be difficult to infect when exposed to the organism; treating or culling infected females; and initiating an immunization program.

Protection of a herd from vibriosis by vaccination has proven to be effective. To induce an immune response to vibriosis, heifers should be vaccinated two or three times at one-month intervals after they are 6 months of age for the primary immunization. If risk of exposure to carrier males or females is present, annual vaccinations should be used to booster immunity.

### Anaplasmosis

Abortion following maternal infection with anaplasmosis is an indirect result of poor oxygen transport to the fetus. The organism is transmitted by blood-carrying vectors such as ticks, blood-sucking flies, mosquitoes and surgical instruments. Control of the disease is dependent on controlling blood-sucking insects, eliminating carrier animals from the herd by test and cull, treatment with tetracycline antibiotics, and in some

cases utilizing vaccines.

Killed vaccines, which provide a low level of resistance to anaplasmosis, are available on an experimental basis. A vaccinated animal is still capable of becoming infected with anaplasmosis and becoming a carrier, but should be protected from symptoms, including abortion. A reaction causing a calf to destroy its own red blood cells has

occurred in calves suckling cows vaccinated with previously available vaccines, so risk assessment should determine the benefits vs. the risks involved.

### *Haemophilus somnus*

*Haemophilus somnus* can cause male and female infertility, and, only rarely,

*(Continued on page 48)*

---

**Vaccination programs should not be the cornerstone for a herd health program, but rather a part of a sound management program.**

---

**The Veterinary Link** (from page 47)

abortion. Transmission for the abortion syndrome is uncertain, but is most likely by mouth. Data showing the ability of vaccination to protect against abortion is lacking. Like most bacterial vaccines, protection from clinical disease is short-lived at best; therefore, a minimum of two initial vaccinations given at one-

month intervals and at least annual boosters would be needed to provide even theoretical protection in those herds where the disease has been demonstrated.

**Trichomoniasis**

Trichomoniasis is caused by a protozoa organism (not a bacteria or virus). The organism is passed from one

animal to the other during the act of mating. In cows, the parasite is confined to the reproductive tract. Trichomonades damage host tissue, cause inflammation of the uterus and invade placental and fetal tissue, resulting in early embryonic death. After a variable period of infertility, cows are usually able to clear the infection (although PI females have been reported). On subsequent exposure

to infected bulls, cows appear to be less susceptible to infection.

In bulls, trichomoniasis does not have any symptoms. The organisms are located on the surfaces of the penis and penile sheath, where they cause little damage. Most bulls less than 4 years of age appear to recover spontaneously or to resist infection.

Control of trichomoniasis outbreaks involves management practices such as using AI, using bulls less than 4 years of age, culling females that do not conceive in a short breeding season, and continuing surveillance of the herd by culturing bulls and culling carriers. Vaccination programs for females exposed to trichomoniasis-infected bulls appear to be beneficial in controlling outbreaks. The program should include two vaccinations four weeks apart for the initial vaccination and annual boosters thereafter. Researchers have shown that, although an immunization program did not completely prevent trichomoniasis, it did decrease the number of cows infected and decreased the length of time it took for infected cows to clear the organism.

Preventing the introduction of trichomoniasis into a herd in areas where the disease is common includes eliminating common pastures and identifying carriers by examination of samples taken from the penile sheath from bulls three times at weekly intervals before the start of the breeding season.

**Neospora caninum**

A vaccine has been licensed for use in the United States to aid in the control of abortion due to neospora, but field trials demonstrating its ability to decrease the risk of infection or abortion are not available. In addition, use of the vaccine may complicate or eliminate the ability to utilize a test-and-cull strategy, because veterinarians may not be able to tell the difference between antibody responses caused by natural infection and vaccination.

**Manage herd health**

Protection from pregnancy wastage in beef cattle has been enhanced by the development and improvement of vaccines against several important organisms causing embryonic loss and abortion. In order to obtain benefit from these vaccines, their use must follow the principles of immunology, and they must be combined with sound management practices that control risk.

Because of the limited number of diseases for which vaccines are available and the inability of vaccines to provide immunity in the presence of nutritional deficiencies, immunosuppression and other management problems, vaccination programs should not be the cornerstone for a herd health program, but rather a part of a sound management program.



**AMERICAN ANGUS ASSOCIATION**  
THE BUSINESS BREED

**APPLICATION FOR REGULAR MEMBERSHIP**

American Angus Association, 3201 Frederick Avenue, Saint Joseph, Missouri 64506  
(816) 383-5100 • Fax: (816) 238-9703 • www.angus.org

I (we) hereby make application to the Board of Directors of the AMERICAN ANGUS ASSOCIATION for Regular Membership and, in consideration of acceptance, agree to conform to and abide by the By-Laws, Rules and Regulations of the Association and Amendments or Modifications thereto which may, from time to time, be adopted. By signing this application for membership, applicant irrevocably waives any claim against and grants an absolute release to the American Angus Association, any member, employee or agent of the Association, for any act or omission in connection with the Association, including but not limited to, any enforcement of the rules and regulations presently in effect or hereafter adopted by the Association. It is requested that the Membership be entered as follows:

**Membership Name** \_\_\_\_\_ **Location of Herd** \_\_\_\_\_  
PLEASE PRINT (Limit Name to 74 Spaces) (City) (County) (State)

**Mail Address** \_\_\_\_\_  
(P.O. Box, or Street) (City) (County) (State) (Zip)

**E-Mail** \_\_\_\_\_ **Date of Birth (If under 21 years of age)** \_\_\_\_\_

**Phone # Home** ( ) \_\_\_\_\_ **Signed** \_\_\_\_\_

**Phone # Business** ( ) \_\_\_\_\_ **Date** \_\_\_\_\_

**Fax #** ( ) \_\_\_\_\_

- Fee of \$80.00 is enclosed and includes a one year subscription to *Angus Journal*
- Certificates enclosed for correction
- Enclosed is my membership fee of \$30.00. I understand this membership does not include subscription to the *Angus Journal*. Initial here \_\_\_\_\_

**Corporations** — In case of a corporation, the corporate name shall be entered on the application, which is to be signed by a duly authorized officer, and below, on this side, at least two officers of the corporation shall sign their names.

**Partnerships** — In case of a firm or partnership, the name of the firm or partnership shall be entered on the application, which is to be signed by one of the partners, and below, on this side, all partners shall sign their names.

**President** \_\_\_\_\_

**Vice Pres.** \_\_\_\_\_

**Secretary** \_\_\_\_\_

**Treasurer** \_\_\_\_\_

**PLEASE READ THE FOLLOWING BEFORE FILLING IN APPLICATION; THEN DETACH**

Regular membership may be applied for in the name of a person, a farm or ranch name; a partnership; or a corporation. Membership fees are payable at the time the application is submitted.

Your regular membership is good for one year. Thereafter, annual dues are required for all regular memberships.

Application should show the name in which the membership is to be entered. If animal registrations are to be applied for in an individual's name the membership should be in the name of the individual; if by a partnership the membership should be in the exact name of the partnership; if by a corporation the membership should be in the exact corporate name. All partners in a partnership, and at least two officers of a corporation must sign their names on the application.

The length of the membership name and location must not exceed 40 spaces, for example:

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40  
 J o n e s & S m i t h C a t t l e C o R o c k R i v e r I A

WHEN SUBMITTING THE APPLICATION PLEASE ENCLOSE THE REGISTRATION CERTIFICATES OF THE ANGUS CATTLE YOU NOW OWN. ASSOCIATION RECORDS ARE KEPT IN MEMBER AND NON-MEMBER SECTION OF THE FILLS, AND THE CERTIFICATES WILL ENABLE US TO REMOVE THEIR RECORD FROM THE NON-MEMBER GROUP AND PLACE THEM UNDER YOUR NAME IN THE MEMBERSHIP SECTION. YOUR REGISTRATION CERTIFICATES WILL BE RE-ISSUED SHOWING YOUR EXACT MEMBERSHIP NAME AND MEMBER CODE NUMBER. THIS WILL GREATLY FACILITATE REGISTRATION OF NEW CALVES WHEN YOU SUBMIT APPLICATIONS OR TRANSFER ANIMALS YOU NOW OWN.